



Adverse health effects of 5G mobile networking technology under real-life conditions

Ronald N. Kostoff^{a,*}, Paul Heroux^b, Michael Aschner^c, Aristides Tsatsakis^{d,e}

^a Research Affiliate, School of Public Policy, Georgia Institute of Technology, Georgia, United States

^b Toxicology and Health Effects of Electromagnetism, McGill University, Canada

^c Molecular Pharmacology, Einstein Center of Toxicology, Albert Einstein College of Medicine, United States

^d Laboratory of Toxicology, Medical School, University of Crete, Voutes, 71409 Heraklion, Crete, Greece

^e Department of Analytical, Toxicology, Pharmaceutical Chemistry and Pharmacognosy, Sechenov University, 119991 Moscow, Russia

GRAPHICAL ABSTRACT



Adverse Health Effects of Wireless Radiation on Humans

Metabolic Disturbance	Reactive Oxygen Species Generation	Genotoxicity and Carcinogenicity	Immunotoxicity and Inflammation	Apoptosis and Necrosis
Discomfort Symptoms	Sensory Disorders	Sleep Disorders	Congenital Abnormalities	Precancerous Conditions
CANCER	NEURODEGENERATION	INFERTILITY	NEUROBEHAVIORAL	CARDIOVASCULAR

ARTICLE INFO

Keywords:

Electromagnetic fields
Wireless radiation
Non-ionizing radiation
Mobile networking technology
5G
Adverse health effects
Toxicology
Toxic stimuli combinations
Synergistic effects
Combined effects
Systemic effects
Real-life simulation

ABSTRACT

This article identifies adverse effects of non-ionizing non-visible radiation (hereafter called wireless radiation) reported in the premier biomedical literature. It emphasizes that most of the laboratory experiments conducted to date are not designed to identify the more severe adverse effects reflective of the real-life operating environment in which wireless radiation systems operate. Many experiments do not include pulsing and modulation of the carrier signal. The vast majority do not account for synergistic adverse effects of other toxic stimuli (such as chemical and biological) acting in concert with the wireless radiation. This article also presents evidence that the nascent 5G mobile networking technology will affect not only the skin and eyes, as commonly believed, but will have adverse systemic effects as well.

1. Introduction

Wireless communications have been expanding globally at an

exponential rate. The latest imbedded version of mobile networking technology is called 4G (fourth generation), and the next version (called 5G- fifth generation) is in the early implementation stage. Neither 4G

* Corresponding author.

E-mail addresses: rkostoff@gmail.com (R.N. Kostoff), toxlab.uoc@gmail.com (A. Tsatsakis).

<https://doi.org/10.1016/j.toxlet.2020.01.020>

Received 12 December 2019; Received in revised form 16 January 2020; Accepted 23 January 2020

Available online 25 January 2020

0378-4274/ © 2020 Elsevier B.V. All rights reserved.

nor 5G have been tested for safety in credible real-life scenarios. Alarming, many of the studies conducted in more benign environments show harmful effects from this radiation. The present article overviews the medical and biological studies that have been performed to date relative to effects from wireless radiation, and shows why these studies are deficient relative to safety. However, even in the absence of the missing real-life components such as toxic chemicals and biotoxins (which tend to exacerbate the adverse effects of the wireless radiation), the literature shows there is much valid reason for concern about potential adverse health effects from both 4G and 5G technology. The studies on wireless radiation health effects reported in the literature should be viewed as extremely conservative, substantially underestimating the adverse impacts of this new technology.

2. Wireless radiation/electromagnetic spectrum

This section overviews the electromagnetic spectrum, and delineates the parts of the spectrum on which this article will focus. The electromagnetic spectrum encompasses the entire span of electromagnetic radiation, including:

- ionizing radiation (gamma rays, x-rays, and the extreme ultraviolet, with wavelengths below $\sim 10^{-7}$ m and frequencies above $\sim 3 \times 10^{15}$ Hz);
- non-ionizing visible radiation (wavelengths from $\sim 4 \times 10^{-7}$ m to $\sim 7 \times 10^{-7}$ m and frequencies between $\sim 4.2 \times 10^{14}$ Hz and $\sim 7.7 \times 10^{14}$ Hz);
- non-ionizing non-visible radiation

short wavelength radio waves and microwaves, with wavelengths between $\sim 10^{-3}$ m and $\sim 10^5$ m and frequencies between $\sim 3 \times 10^{11}$ to $\sim 3 \times 10^3$ Hz;

long wavelengths, ranging between $\sim 10^5$ m and $\sim 10^8$ m and frequencies ranging between 3×10^3 and 3 Hz.

How are these frequencies used in practice?

- The low frequencies (3 Hz–300 KHz) are used for electrical power line transmission (60 Hz in the U.S.) as well as maritime and submarine navigation and communications.
- Medium frequencies (300 KHz–900 MHz) are used for AM/FM/TV broadcasts in North America.
- Lower microwave frequencies (900 MHz–5 GHz) are used for telecommunications such as microwave devices/communications, radio astronomy, mobile/cell phones, and wireless LANs.
- Higher microwave frequencies (5 GHz–300 GHz) are used for radar and proposed for microwave WiFi, and will be used for high-performance 5 G.
- Terahertz frequencies (300 GHz–3000 GHz) are used increasingly for imaging to supplement X-rays in some medical and security scanning applications (Kostoff and Lau, 2017).

In the present study of wireless radiation health effects, the frequency spectrum ranging from 3 Hz to 300 GHz is covered, with particular emphasis on the high frequency communications component ranging from ~ 1 GHz to ~ 300 GHz. Why was this part of the spectrum selected? Previous reviews of wireless radiation health effects found that pulsed electromagnetic fields (PEMF) applied for relatively short periods of time could sometimes be used for therapeutic purposes, whereas chronic exposure to electromagnetic fields (EMF) in the power frequency range (~ 60 Hz) and microwave frequency range (~ 1 GHz–tens GHz) tended to result in detrimental health effects (Kostoff and Lau, 2013, 2017). Given present concerns about the rapid expansion of 5G communications systems (which are projected to use mainly the higher microwave frequencies part of the spectrum in the highest performance (aka high-band) mode) in the absence of adequate and rigorous safety testing, more emphasis will be placed on the

communications frequencies in this document.

3. Modern wireless radiation exposures

In ancient times, sunlight and its lunar reflections provided the bulk of the visible spectrum for human beings (with fire a distant second and lightning a more distant third). Now, many varieties of artificial light (incandescent, fluorescent, and light emitting diode) have replaced the sun as the main supplier of visible radiation during waking hours. Additionally, EMF radiations from other parts of the non-ionizing non-visible spectrum have become ubiquitous in daily life, such as from wireless computing and telecommunications. In the last two or three decades, the explosive growth in the cellular telephone industry has placed many residences in metropolitan areas within less than a mile of a cell tower. Future implementation of the next generation of mobile networking technology, 5 G, will increase the cell tower densities by an order of magnitude. Health concerns have been raised about wireless radiation from (1) mobile communication devices, (2) occupational exposure, (3) residential exposure, (4) wireless networks in homes, businesses, and schools, (5) automotive radar, and (6) other non-ionizing EMF radiation sources, such as ‘smart meters’ and ‘Internet of Things’.

4. Demonstrated biological and health effects from prior generations of wireless networking technology

There have been two major types of studies performed to ascertain biological and health effects of wireless radiation: laboratory and epidemiology. The laboratory tests performed provided the best scientific understanding of the effects of wireless radiation, but did not reflect the real-life environment in which wireless radiation systems operate (exposure to toxic chemicals, biotoxins, other forms of toxic radiation, etc). There are three main reasons the laboratory tests failed to reflect real-life exposure conditions for human beings.

First, the laboratory tests have been performed mainly on animals, especially rats and mice. Because of physiological differences between small animals and human beings, there have been continual concerns about extrapolating small animal results to human beings. Additionally, while inhaled or ingested substances can be scaled from laboratory experiments on small animals to human beings relatively straight-forwardly, radiation may be more problematic. For non-ionizing radiation, penetration depth is a function of frequency, tissue, and other parameters. Radiation could penetrate much deeper into a small animal’s interior than similar wavelength radiation in humans, because of the much smaller animal size. Different organs and tissues would be affected, with different levels of power density.

Second, the typical incoming EMF signal for many/most laboratory tests performed in the past consisted of single carrier wave frequency; the lower frequency superimposed signal containing the information was not always included. This omission may be important. As Panagopoulos states: “It is important to note that except for the RF/microwave carrier frequency, Extremely Low Frequencies - ELF (0–3000 Hz) are always present in all telecommunication EMFs in the form of pulsing and modulation. There is significant evidence indicating that the effects of telecommunication EMFs on living organisms are mainly due to the included ELF’s... While ~ 50 % of the studies employing simulated exposures do not find any effects, studies employing real-life exposures from commercially available devices display an almost 100 % consistency in showing adverse effects” (Panagopoulos, 2019). These effects may be exacerbated further with 5 G: “with every new generation of telecommunication devices.....the amount of information transmitted each moment.....is increased, resulting in higher variability and complexity of the signals with the living cells/ organisms even more unable to adapt” (Panagopoulos, 2019).

Third, these laboratory experiments typically involved one stressor

(toxic stimulus) and were performed under pristine conditions. This contradicts real-life exposures, where humans are exposed to multiple toxic stimuli, in parallel or over time (Tsatsakis et al., 2016, 2017; Docea et al., 2019a). In perhaps five percent of the cases reported in the wireless radiation literature, a second stressor (mainly a biological or chemical toxic stimulus) was added to the wireless radiation stressor, to ascertain whether additive, synergistic, potentiative, or antagonistic effects were generated by the combination (Kostoff and Lau, 2013, 2017; Juutilainen, 2008; Juutilainen et al., 2006).

Combination experiments are extremely important because, when other toxic stimuli are considered in combination either with each other or with wireless radiation, the synergies tend to enhance the adverse effects of each stimulus in isolation. This was shown in several studies that evaluated the cumulative effects of chronic exposure to low doses of xenobiotics in combination (Kostoff et al., 2018; Docea et al., 2018; Tsatsakis et al., 2019a; Docea et al., 2019b; Tsatsakis et al., 2019b, c; Fountoucidou et al., 2019). For those combinations that include wireless radiation, combined exposure to toxic stimuli and wireless radiation translates into much lower levels of tolerance for each toxic stimulus in the combination relative to its exposure levels that produce adverse effects in isolation. Accordingly, the exposure limits for wireless radiation when examined in combination with other potentially toxic stimuli would be far lower for safety purposes than those derived from wireless radiation exposures in isolation.

Thus, almost all of the wireless radiation laboratory experiments that have been performed to date are flawed/limited with respect to showing the full adverse impact of the wireless radiation that would be expected under real-life conditions. Either 1) non-inclusion of signal information or 2) using single stressors only tends to underestimate the seriousness of the adverse effects from wireless radiation. Excluding **both** of these phenomena from experiments, as was done in the vast majority of the reported wireless radiation health effects studies, tends to amplify this underestimation substantially. Thus, the results reported in the biomedical literature should be viewed as 1) extremely conservative and 2) the very low ‘floor’ of the seriousness of the adverse effects from wireless radiation, not the ‘ceiling’.

In contrast to the controlled pristine environments that characterize the wireless radiation animal laboratory experiments, the wireless radiation epidemiology studies carried out to date typically involved human beings who had been subjected to myriad known and unknown stressors prior to (and during) the study. The real-life human exposure levels from cell tower studies (reported by Kostoff and Lau (2017)) that showed increased cancer incidence were orders of magnitude lower than those exposure levels generated in the recent highly-funded National Toxicology Program animal laboratory studies (Melnick, 2019). We believe the inclusion of real-world effects in the cell tower studies accounted for the orders of magnitude exposure level decreases that were associated with the increased cancer incidence. The laboratory tests were conducted under controlled conditions not reflective of real-life, while the epidemiology studies were performed in the presence of many stressors, known and unknown, reflective of real-life. The myriad toxic stimuli exposure levels of the epidemiology studies were, for the most part, uncontrolled.

A vast literature published over the past sixty years shows adverse effects from wireless radiation applied in isolation or as part of a combination with other toxic stimuli. Extensive reviews of wireless radiation-induced biological and health effects have been published (Kostoff and Lau, 2013, 2017; Belpomme et al., 2018; Desai et al., 2009; Di Ciaula, 2018; Doyon and Johansson, 2017; Havas, 2017; Kaplan et al., 2016; Lerchl et al., 2015; Levitt and Lai, 2010; Miller et al., 2019; Pall, 2016, 2018; Panagopoulos, 2019; Panagopoulos et al., 2015; Russell, 2018; Sage and Burgio, 2018; van Rongen et al., 2009; Yakymenko et al., 2016; Bioinitiative, 2012). In aggregate, for the high frequency (radiofrequency-RF) part of the spectrum, these reviews show that RF radiation **below the FCC guidelines** can result in:

- carcinogenicity (brain tumors/glioma, breast cancer, acoustic neuromas, leukemia, parotid gland tumors),
- genotoxicity (DNA damage, DNA repair inhibition, chromatin structure),
- mutagenicity, teratogenicity,
- neurodegenerative diseases (Alzheimer’s Disease, Amyotrophic Lateral Sclerosis),
- neurobehavioral problems, autism, reproductive problems, pregnancy outcomes, excessive reactive oxygen species/oxidative stress, inflammation, apoptosis, blood-brain barrier disruption, pineal gland/melatonin production, sleep disturbance, headache, irritability, fatigue, concentration difficulties, depression, dizziness, tinnitus, burning and flushed skin, digestive disturbance, tremor, cardiac irregularities,
- adverse impacts on the neural, circulatory, immune, endocrine, and skeletal systems.

From this perspective, RF is a highly pervasive cause of disease!

The response from industry has been that no mechanism could explain the biological action of non-thermal and non-ionizing EM fields. Yet, reports of clear perturbations of biological systems at levels near or even below 1000 $\mu\text{W}/\text{m}^2$ (Bioinitiative, 2019) were explained by perturbations in electron and proton transfers supporting ATP production in mitochondria (Sanders et al., 1980; 1985) exposed to RF or ELF signals (Li and Heroux, 2014).

To obtain another perspective on the full spectrum of adverse effects from wireless radiation, a query was run on Medline to retrieve representative records associated with adverse EMF effects (mainly, but not solely, RF). Over 5400 records were retrieved, and the leading Medical Subject Headings (MeSH) extracted. The categories of adverse impacts from both approaches match quite well. The adverse health effects range from myriad feelings of discomfort to life-threatening diseases.

The full list of MeSH Headings associated with this retrieval is shown in Appendix 1 of (Kostoff, 2019). The interested reader can ascertain what other diseases/symptoms were included. The 5400+ references retrieved are shown in Appendix 2 of (Kostoff, 2019).

5. What types of biological and health effects can be expected from 5G wireless networking technology?

The potential 5G adverse effects derive from the intrinsic nature of the radiation, and its interaction with tissue and target structures. 4G networking technology was associated mainly with carrier frequencies in the range of $\sim 1\text{--}2.5$ GHz (cell phones, WiFi). The wavelength of 1 GHz radiation is 30 cm, and the penetration depth in human tissue is a few centimeters. In its highest performance (high-band) mode, 5G networking technology is mainly associated with carrier frequencies at least an order of magnitude greater than the 4G frequencies, although, as stated previously, “ELFs (0–3000 Hz) are always present in all telecommunication EMFs in the form of pulsing and modulation”. Penetration depths for the carrier frequency component of high-band 5G wireless radiation will be on the order of a few millimeters (Alekseev et al., 2008a, b). At these wavelengths, one can expect resonance phenomena with small-scale human structures (Betzalel et al., 2018). Additionally, numerical simulations of millimeter-wave radiation resonances with insects showed a general increase in absorbed RF power at and above 6 GHz, in comparison to the absorbed RF power below 6 GHz. A shift of 10 % of the incident power density to frequencies above 6 GHz was predicted to lead to an **increase in absorbed power between 3–370 %** (Thielens et al., 2018).

The common ‘wisdom’ presented in the literature and media is that, if there are adverse impacts resulting from high-band 5G, the main impacts will be focused on near-surface phenomena, such as skin cancer, cataracts, and other skin conditions. However, there is evidence that biological responses to millimeter-wave irradiation can be initiated

within the skin, and the subsequent systemic signaling in the skin can result in physiological effects on the nervous system, heart, and immune system (Russell, 2018).

Additionally, consider the following reference (Zalyubovskaya, 1977). This is one of many translations of articles produced in the Former Soviet Union on wireless radiation (also, see reviews of Soviet research on this topic by McRee (1979, 1980), Kositsky et al. (2001), and Glaser and Dodge (1976)). On p. 57 of the pdf link, the article by Zalyubovskaya addresses biological effects of millimeter radiowaves. Zalyubovskaya ran experiments using power fluxes of 10,000,000 $\mu\text{W}/\text{square meter}$ (the FCC (Federal Communications Commission) guideline limit for the general public today in the USA), and frequencies on the order of 60 GHz. Not only was skin impacted adversely, but also heart, liver, kidney, spleen tissue as well, and blood and bone marrow properties. These results reinforce the conclusion of Russel (quoted above) that **systemic results may occur from millimeter-wave radiation**. To re-emphasize, for Zalyubovskaya's experiments, the incoming signal was unmodulated carrier frequency only, and the experiment was single stressor only. Thus, the expected real-world results (when human beings are impacted, the signals are pulsed and modulated, and there is exposure to many toxic stimuli) would be far more serious and would be initiated at lower (perhaps much lower) wireless radiation power fluxes.

The Zalyubovskaya paper was published in 1977. The referenced version was classified in 1977 by USA authorities and declassified in 2012. What national security concerns caused it (and the other papers in the linked pdf reference) to be classified for 35 years, until declassification in 2012? Other papers on this topic with similar findings were published in the USSR (and the USA) at that time, or even earlier, but many never saw the light of day, both in the USSR and the USA. It appears that the potentially damaging effects of millimeter-wave radiation on the skin (and other major systems in the body) have been recognized for well over forty years, yet today's discourse only revolves around the possibility of modest potential effects on the skin and perhaps cataracts from millimeter-wave wireless radiation.

6. What is the consensus on adverse effects from wireless radiation?

Not all studies of wireless radiation have shown adverse effects. For example, consider potential genotoxic effects of mobile phone radiation. A study investigating "the effect of mobile phone use on genomic instability of the human oral cavity's mucosa cells" concluded "Mobile phone use did not lead to a significantly increased frequency of micronuclei" (Hintzsche and Stopper, 2010).

Conversely, a 2017 study investigated buccal cell preparations for genomic instability, and found "The frequency of micronuclei (13.66x), nuclear buds (2.57x), basal (1.34x), karyorrhectic (1.26x), karyolytic (2.44x), pyknotic (1.77x) and condensed chromatin (2.08x) cells were highly significantly ($p = 0.000$) increased in mobile phone users" (Gandhi et al., 2017). Also, a 2017 study to ascertain the "effect of cell phone emitted radiations on the orofacial structures" concluded that "Cell phone emitted radiation causes nuclear abnormalities of the oral mucosal cells" (Mishra et al., 2017). Further, a 2016 study to "explore the effects of mobile phone radiation on the MN frequency in oral mucosal cells" concluded "The number of micronucleated cells/1000 exfoliated buccal mucosal cells was found to be significantly increased in high mobile phone users group than the low mobile phone users group" (Banerjee et al., 2016). Finally, a study aimed at investigating the health effects of WiFi exposure concluded "long term exposure to WiFi may lead to adverse effects such as neurodegenerative diseases as observed by a **significant alteration on AChE gene expression** and some neurobehavioral parameters associated with brain damage" (Obajuluwa et al., 2017).

There are many possible reasons to explain this lack of consensus.

- 1) There may be 'windows' in parameter space where adverse effects occur, and operation outside these windows would show a) no effects or b) hormetic effects or c) therapeutic effects. For example, if information content of the signal is a strong contributor to adverse health effects (Panagopoulos, 2019), then experiments that involve only the carrier frequencies may be outside the window where adverse health effects occur. Alternatively, in this specific example, the carrier signal and the information signal could be viewed as a combination of potentially toxic stimuli, where the adverse effects of each component are enabled because of the synergistic effects of the combination.

As another example, an adverse health impact on one strain of rodent was shown for a combination of 50 Hz EMF and DMBA, while no adverse health impact was shown on another rodent strain for the same toxic stimuli combination (Fedrowitz et al., 2004). From a higher-order combination perspective, if genetic abnormalities/differences are viewed conceptually as potentially equivalent to a toxic stimulus for combination purposes, then a synergistic three-constituent combination of 50 Hz EMF, DMBA, and genetics was required to produce adverse health impacts in the above experiment. If these results can be extrapolated across species, then human beings could exhibit different responses to the same electromagnetic stimuli based on their unique genetic predispositions (Caccamo et al., 2013; De Luca et al., 2014).

- 1) Research quality could be poor, and adverse effects were overlooked.
- 2) Or, the research team could have had a preconceived agenda, where finding no adverse effects from wireless radiation was **THE** objective of the study. For example, studies have shown that industry-funded research of wireless radiation adverse health effects is far more likely to show no effects than funding from non-industry sources (Huss et al., 2007; Slesin, 2006; Carpenter, 2019). Studies in disciplines other than wireless radiation have shown that, for products of high military, commercial, and political sensitivity, 'researchers'/organizations are hired to publish articles that conflict with the credible science, and therefore create doubt as to whether the product of interest is harmful (Michaels, 2008; Oreskes and Conway, 2011). Unfortunately, given the strong dependence of the civilian and military economies on wireless radiation, incentives for identifying adverse health effects from wireless radiation are minimal and disincentives are many. These perverse incentives apply not only to the sponsors of research and development, but to the performers as well.

Even the Gold Standard for research credibility - **independent replication of research results** - is questionable in politically, commercially, and militarily sensitive areas like wireless radiation safety, where the accelerated implementation goals of most wireless radiation research sponsors (government and industry) are aligned. It is imperative that highly objective evaluators with minimal conflicts of interest play a central role ensuring that rigorous safety standards for wireless radiation systems are met before widescale implementation is allowed.

7. Conclusions

Wireless radiation offers the promise of improved remote sensing, improved communications and data transfer, and improved connectivity. Unfortunately, there is a large body of data from laboratory and epidemiological studies showing that previous and present generations of wireless networking technology have significant adverse health impacts. Much of this data was obtained under conditions not reflective of real-life. When real-life considerations are added, such as 1) including the information content of signals along with 2) the carrier frequencies, and 3) including other toxic stimuli in combination with

the wireless radiation, the adverse effects associated with wireless radiation are increased substantially. Superimposing 5G radiation on an already imbedded toxic wireless radiation environment will exacerbate the adverse health effects shown to exist. Far more research and testing of potential 5G health effects under real-life conditions is required before further rollout can be justified.

Transparency document

The [Transparency document](#) associated with this article can be found in the online version.

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

References

Alekseev, S.I., Radziewsky, A.A., Logani, M.K., Ziskin, M.C., 2008a. Millimeter wave dosimetry of human skin. *Bioelectromagnetics* 29 (January (1)), 65–70. <https://doi.org/10.1002/bem.20363>.

Alekseev, S.I., Gordienko, O.V., Ziskin, M.C., 2008b. Reflection and penetration depth of millimeter waves in murine skin. *Bioelectromagnetics* 29 (July (5)), 340–344. <https://doi.org/10.1002/bem.20401>.

Banerjee, S., Singh, N.N., Sreedhar, G., Mukherjee, S., 2016. Analysis of the genotoxic effects of mobile phone radiation using buccal micronucleus assay: a comparative evaluation. *J. Clin. Diagn. Res.* 10 (3) ZC82-ZC85.

Belpomme, D., Hardell, L., Belyaev, I., Burgio, E., Carpenter, D.O., 2018. Thermal and non-thermal health effects of low intensity non-ionizing radiation: an international perspective. *Environ. Pollut.* 242, 643–658.

Betzalel, N., Ben Ishai, P., Feldmann, Y., 2018. The human skin as a sub-THz receiver - Does 5G pose a danger to it or not? *Environ. Res.* 163, 208–216.

BioInitiative Working Group, Cindy Sage, David O. Carpenter, BioInitiative Report: A Rationale for Biologically-Based Public Exposure Standards for Electromagnetic Radiation at 2012; www.bioinitiative.org December 31, 2012, last updated 2019.

Caccamo, D., Cesareo, E., Mariani, S., Raskovic, D., Lentile, R., Currò, M., Korkina, L., De Luca, C., 2013. Xenobiotic sensor- and metabolism-related gene variants in environmental sensitivity-related illnesses: a survey on the Italian population. *Oxid. Med. Cell. Longev.* 2013 <https://doi.org/10.1155/2013/831969>. Article ID 831969.

Carpenter, D.O., 2019. Extremely low frequency electromagnetic fields and cancer: how source of funding affects results. *Environ. Res.* 178, 108688. <https://doi.org/10.1016/j.envres.2019.108688>.

De Luca, C., Thai, J.C., Raskovic, D., Cesareo, E., Caccamo, D., Trukhanov, A., Korkina, L., 2014. Metabolic and genetic screening of electromagnetic hypersensitive subjects as a feasible tool for diagnostics and intervention. *Mediators Inflamm.* 2014, 924184. <https://doi.org/10.1155/2014/924184>. Epub 2014 Apr 9.

Desai, N.R., Kesari, K.K., Agarwal, A., 2009. Pathophysiology of cell phone radiation: oxidative stress and carcinogenesis with focus on male reproductive system. *Reprod. Biol. Endocrinol.* 7.

Di Ciaula, A., 2018. Towards 5G communication systems: are there health implications? *Int. J. Hyg. Environ. Health* 221 (3), 367–375.

Docea, A.O., Gofita, E., Goumenou, M., Calina, D., Rogoveanu, O., Varut, M., Olaru, C., Kerasioti, E., Fountoucidou, P., Taitzoglou, I., Zlatian, O., Rakitskii, V.N., Hernandez, A.F., Kouretas, D., Tsatsakis, A., 2018. Six months exposure to a real life mixture of 13 chemicals' below individual NOAELs induced non monotonic sex-dependent biochemical and redox status changes in rats. *Food Chem. Toxicol.* 115, 470–481 [doi: 10.1016/j.fct.2018.03.052](https://doi.org/10.1016/j.fct.2018.03.052).

Docea, A.O., Calina, D., Goumenou, M., Neagu, M., Gofita, Tsatsakis, A., 2019a. Study design for the determination of toxicity from long-term-low-dose exposure to complex mixtures of pesticides, food additives and lifestyle products. *Toxicol. Lett.* 258 <https://doi.org/10.1016/j.toxlet.2016.06.1666>. S179-S179.

Docea, A.O., Goumenou, M., Calina, D., Arsene, A.L., Dragoi, C.M., Gofita, E., Pisoschi, C.G., Zlatian, O., Stivaktakis, P.D., Nikolouzakis, T.K., Kalođeraki, A., Izotov, B.N., Galateanu, B., Hudita, A., Calabrese, E.J., Tsatsakis, A., 2019b. Adverse and hormetic effects in rats exposed for 12 months to low dose mixture of 13 chemicals: RLR5 part III. *Toxicol. Lett.* 310, 70–91. <https://doi.org/10.1016/j.toxlet.2019b.04.005>.

Doyon, P.R., Johansson, O., 2017. Electromagnetic fields may act via calcineurin inhibition to suppress immunity, thereby increasing risk for opportunistic infection: conceivable mechanisms of action. *Med. Hypotheses* 106, 71–87.

Fedrowitz, M., Kamino, K., Loscher, W., 2004. Significant differences in the effects of magnetic field exposure on 7,12-dimethylbenz(a)anthracene-induced mammary carcinogenesis in two substrains of Sprague-Dawley rats. *Cancer Res.* 64 (1), 243–251.

Fountoucidou, P., Veskoukis, A.S., Kerasioti, E., Docea, A.O., Taitzoglou, I.A., Liesivuori, J., Tsatsakis, A., 2019. Kouras DA mixture of routinely encountered xenobiotics induces both redox adaptations and perturbations in blood and tissues of rats after a long-term low-dose exposure regimen: the time and dose issue. *Toxicol. Lett.* 317, 24–44. <https://doi.org/10.1016/j.toxlet.2019.09.015>.

Gandhi, G., Singh, P., Kaur, G., 2017. Perspectives revisited - the buccal cytochrome assay in mobile phone users. *Int. J. Hum. Genet.* 15 (4), 173–182.

Glaser, R., Dodge, C.H., 1976. Biomedical Aspects of Radiofrequency Radiation: A Review of Selected Soviet, East European, and Western References. Selected Papers of the USNC/URSI Annuit Meeting (Boulder, CO. 1975). HEW Publication (FDA) 77-8010/8011, pp. 2–34 Dec.

Havas, M., 2017. When theory and observation collide: can non-ionizing radiation cause cancer? *Environ. Pollut.* 221, 501–505.

Hintzsche, H., Stopper, H., 2010. Micronucleus frequency in buccal mucosa cells of mobile phone users. *Toxicol. Lett.* 193 (1), 124–130.

Huss, A., Egger, M., Hug, K., Huwiler-Müntener, K., Rössli, M., 2007. Source of funding and results of studies of health effects of mobile phone use: systematic review of experimental studies. *Environ. Health Perspect.* 115 (1), 1–4.

Juutilainen, J., 2008. Do electromagnetic fields enhance the effects of environmental carcinogens? *Radiat. Prot. Dosimetry* 132 (2), 228–231.

Juutilainen, J., Kumlin, T., Naarala, J., 2006. Do extremely low frequency magnetic fields enhance the effects of environmental carcinogens? A meta-analysis of experimental studies. *Int. J. Radiat. Biol.* 82 (1), 1–12.

Kaplan, S., Deniz, O.G., Onger, M.E., Turkmen, A.P., Yurt, K.K., Aydin, I., et al., 2016. Electromagnetic field and brain development. *J. Chem. Neuroanat.* 75, 52–61.

Kositsky, N.N., Nizhelska, A.I., Ponezh, G.V., 2001. Influence of high-frequency electromagnetic radiation at non-thermal intensities on the human body (a review of work by Russian and Ukrainian researchers). No Place to Hide- Newsletter of the Cellular Phone Taskforce Inc. 3 (Supplement (1)).

Kostoff, R.N., 2019. Adverse Effects of Wireless Radiation. PDF. <http://hdl.handle.net/1853/61946>.

Kostoff, R.N., Lau, C.G.Y., 2013. Combined biological and health effects of electromagnetic fields and other agents in the published literature. *Technol. Forecast. Soc. Change* 80 (7), 1331–1349.

Kostoff, R.N., Lau, C.G.Y., 2017. Modified health effects of non-ionizing electromagnetic radiation combined with other agents reported in the biomedical literature. In: Geddes, C.D. (Ed.), *Microwave Effects on DNA and Proteins*. Springer International Publishing AG. https://doi.org/10.1007/978-3-319-50289-2_4.

Kostoff, R.N., Goumenou, M., Tsatsakis, A., 2018. The role of toxic stimuli combinations in determining safe exposure limits. *Toxicol. Rep.* 5, 1169–1172. <https://doi.org/10.1016/j.toxrep.2018.10.010>. eCollection 2018.

Lerchl, A., Klose, M., Grote, K., Wilhelm, A.F.X., Spathmann, O., Fiedler, T., et al., 2015. Tumor promotion by exposure to radiofrequency electromagnetic fields below exposure limits for humans. *Biochem. Biophys. Res. Commun.* 459 (4), 585–590.

Levitt, B.B., Lai, H., 2010. Biological effects from exposure to electromagnetic radiation emitted by cell tower base stations and other antenna arrays. *Environ. Res.* 118, 369–395.

Li, Y., Héroux, P., 2014. Extra-Low-Frequency magnetic fields alter cancer cells through metabolic restriction. *Electromagn. Biol. Med.* 33 (4), 264–275. <https://doi.org/10.3109/15368378.2013.817334>.

McRee, D.I., 1979. Review of Soviet Eastern-European research on health-aspects of microwave-radiation. *Bull. N. Y. Acad. Med.* 55 (11), 1133–1151.

McRee, D.I., 1980. Soviet and Eastern-European research on biological effects of microwave-radiation. *Proc. IEEE* 68 (1), 84–91. <https://doi.org/10.1109/PROC.1980.11586>. Also. https://www.avaate.org/IMG/pdf/mcree80_rev_soviet.pdf.

Melnick, R.L., 2019. Commentary on the utility of the National Toxicology Program study on cell phone radiofrequency radiation data for assessing human health risks despite unfounded criticisms aimed at minimizing the findings of adverse health effects. *Environ. Res.* 168, 1–6. <https://doi.org/10.1016/j.envres.2018.09.010>.

Michaels, D., 2008. *Doubt Is Their Product: How Industry's Assault on Science Threatens Your Health*, 1st edition. Oxford University Press, Oxford, United Kingdom.

Miller, A.B., Sears, M.E., Morgan, L.L., Davis, D.L., Hardell, L., Oremus, M., et al., 2019. Risks to health and well-being from radio-frequency radiation emitted by cell phones and other wireless devices. *Front. Public Health* 7, 223. <https://doi.org/10.3389/fpubh.2019.00223>.

Mishra, S.K., Chowdhary, R., Kumari, S., Rao, S.B., 2017. Effect of cell phone radiations on orofacial structures: a systematic review. *J. Clin. Diagn. Res.* 11 (5), 5.

Obajuluwa, A.O., Akinyemi, A.J., Afolabi, O.B., Adekoya, K., Sanya, J.O., Ishola, A.O., 2017. Exposure to radio-frequency electromagnetic waves alters acetylcholinesterase gene expression, exploratory and motor coordination-linked behaviour in male rats. *Toxicol. Rep.* 4, 530–534.

Oreskes, N., Conway, E.M., 2011. *Merchants of Doubt: How a Handful of Scientists Obscured the Truth on Issues From Tobacco Smoke to Global Warming*. Bloomsbury Press, New York, NY.

Pall, M.L., 2016. Microwave frequency electromagnetic fields (EMFs) produce widespread neuropsychiatric effects including depression. *J. Chem. Neuroanat.* 75, 43–51.

Pall, M.L., 2018. Wi-Fi is an important threat to human health. *Environ. Res.* 164, 405–416.

Panagopoulos, D.J., 2019. Comparing DNA damage induced by mobile telephony and other types of man-made electromagnetic fields. *Mutat. Res. Mutat. Res.* 781, 53–62.

Panagopoulos, D.J., Johansson, O., Carlo, G.L., 2015. Real versus simulated mobile phone exposures in experimental studies. *Biomed Res. Int.* 2015, 607053. <https://doi.org/10.1155/2015/607053>.

Russell, C.L., 2018. 5 G wireless telecommunications expansion: public health and environmental implications. *Environ. Res.* 165, 484–495.

Sage, C., Burgio, E., 2018. Electromagnetic fields, pulsed radiofrequency radiation, and epigenetics: how wireless technologies may affect childhood development. *Child Dev.* 89 (1), 129–136.

Sanders, A.P., Schaefer, D.J., Joines, W.T., 1980. Microwave effects on energy metabolism of rat brain. *Bioelectromagnetics* 1, 171–181.

Sanders, A.P., Joines, W.T., Allis, J.W., 1985. Effects of continuous-wave, pulsed, and

- sinusoidal-amplitude-modulated microwaves on brain energy metabolism. *Bioelectromagnetics*. 6, 89–97.
- Slesin, L., 2006. "Radiation Research" and the Cult of Negative Results. *Microwave News* 31 July. <http://microwavenews.com/RR.html>.
- Thielens, A., Bell, D., Mortimore, D.B., Greco, M.K., Martens, L., Joseph, W., 2018. Exposure of insects to radio-frequency electromagnetic fields from 2 to 120 GHz. *Sci. Rep.* 8, 3924. <https://doi.org/10.1038/s41598-018-22271-3>.
- Tsatsakis, A.M., Docea, A.O., Tsitsimpikou, C., 2016. New challenges in risk assessment of chemicals when simulating real exposure scenarios; simultaneous multi-chemicals' low dose exposure. *Food Chem. Toxicol.* 96, 174–176. <https://doi.org/10.1016/j.fct.2016.08.011>.
- Tsatsakis, A.M., Kouretas, D., Tzatzarakis, M.N., Stivaktakis, P., Tsarouhas, K., Golokhvast, K.S., Rakitskii, V.N., Tutelyan, V.A., Hernandez, A.F., Rezaee, R., Chung, G., Fenga, C., Engin, A.B., Neagu, M., Arsene, A.L., Docea, A.O., Gofita, E., Calina, D., Taitzoglou, I., Liesivuori, J., Hayes, A.W., Gutnikov, S., Tsitsimpikou, C., 2017. Simulating real-life exposures to uncover possible risks to human health: a proposed consensus for a novel methodological approach. *Hum. Exp. Toxicol.* 36 (6), 554–564. <https://doi.org/10.1177/0960327116681652>.
- Tsatsakis, A.M., Docea, A.O., Calina, D., Buga, A.M., Zlatian, O., Gutnikov, S., Kostoff, R.N., Aschner, M., 2019a. Hormetic Neurobehavioral effects of low dose toxic chemical mixtures in real-life risk simulation (RLRS) in rats. *Food Chem. Toxicol.* 125, 141–149. <https://doi.org/10.1016/j.fct.2018.12.043>.
- Tsatsakis, A., Tyshko, N.V., Docea, A.O., Shestakova, S.I., Sidorova, Y.S., Petrov, N.A., Zlatian, O., Mach, M., Hartung, T., Tutelyan, V.A., 2019b. The effect of chronic vitamin deficiency and long term very low dose exposure to 6 pesticides mixture on neurological outcomes - A real-life risk simulation approach. *Toxicol. Lett.* 315, 96–106. <https://doi.org/10.1016/j.toxlet.2019.07.026>.
- Tsatsakis, A., Docea, A.O., Constantin, C., Calina, D., Zlatian, O., Nikolouzakis, T.K., Stivaktakis, P.D., Kalogeraki, A., Liesivuori, J., Tzanakakis, G., Neagu, M., 2019c. Genotoxic, cytotoxic, and cytopathological effects in rats exposed for 18 months to a mixture of 13 chemicals in doses below NOAEL levels. *Toxicol. Lett.* 316, 154–170. <https://doi.org/10.1016/j.toxlet.2019.09.004>.
- van Rongen, E., Croft, R., Juutilainen, J., Lagroye, I., Miyakoshi, J., Saunders, R., et al., 2009. Effects of radiofrequency electromagnetic fields on the human nervous system. *J. Toxicol. Environ. Health-Part B-Crit. Rev.* 12 (8), 572–597.
- Yakymenko, I., Tsybulin, O., Sidorik, E., Henshel, D., Kyrylenko, O., Kyrylenko, S., 2016. Oxidative mechanisms of biological activity of low-intensity radiofrequency radiation. *Electromagn. Biol. Med.* 35 (2), 186–202.
- Zalyubovskaya, N.P., 1977. Biological Effects of Millimeter Radiowaves. *Vrachebnoye Delo*. No. 3. P. 57. <https://www.cia.gov/library/readingroom/docs/CIA-RDP88B01125R000300120005-6.pdf>.